This Page Is Inserted by IFW Operations and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents will not correct images, please do not report the images to the Image Problem Mailbox.

Select a targeted foraging area (i.e., a foraging area to be preconditioned) to be encompassed by the wild ungulate foraging ("WUF") distribution. Select a target period during which the wild ungulates will use the targeted foraging area once it has been preconditioned. Select a preconditioning period during which domesticated ungulates precondition the targeted foraging area prior to the target period. Supply an ingestible in one or more deployment locations within the targeted foraging area, wherein the ingestible is ingestible by domestic ungulates, attracts domestic ungulates to the deployment locations, and increases the dwell time of the ungulates within the targeted foraging area. Allow domesticated ungulates to forage the targeted foraging area (i.e., precondition the targeted foraging area). Remove the domesticated ungulates from the preconditioned targeted foraging area. During the target period, allowing the WUF distribution to shift to encompass the preconditioned targeted foraging area.

F16.1















